

**ORIGINAL RESEARCH ARTICLE****Performance and carcass characteristics of broiler finishers fed diets supplemented with Roxazyme G and exogenous phytase**

***Abu, O. A, Sodeinde, A. O, Ehalodu, J. O, Olomola, O. O, and Babayemi, O. J.**

Department of Animal Science, University of Ibadan, Nigeria

**Corresponding author Email: ohiahmed@yahoo.com*

ABSTRACT

A 21-day feeding trial involving 480, 4-week old unsexed Abhor Acre broilers was carried out in a completely randomised design to evaluate performance, carcass characteristics and cost benefit of broiler finishers fed iso-caloric and iso-nitrogenous corn-soyabean based enzyme supplemented diets. Diet 1 had no supplemented Roxazyme G and phytase (T1-basal diet). The other three diets are: T2-Roxazyme G (0.2g/kg), T3-basal diet +Phytase (1g/kg) and T4-basal diet +Roxazyme G and Phytase (0.2g/kg and 1g/kg) inclusion. The objective was to observe whether addition exogenous enzymes individually or as a mixture improved efficiency of broiler finisher production. The results showed no significant differences ($p>0.05$) across the dietary treatments for daily feed intake, weight gain and FCR. The values obtained were in the range of 137.9g, 1191.9g and 1.45 respectively. There were no significant difference ($p>0.05$) in carcass weight of birds on T4, T2 and T1 but birds fed both Roxazyme G and phytase had heaviest carcass weight of 1533.3g. However, birds fed diets supplemented with phytase alone had significantly lower carcass weight and dressing yield to other dietary treatments. The cut up parts were not significantly different ($p>0.05$) except the weights of the head, drumstick and breast meat. The cost analysis showed that it is not profitable to include the enzymes in the practical broiler finishers' diet used in this study. There was no need to include the above enzymes in the diet of broilers at finishing phase as the birds at this stage can efficiently utilize nutrients in their feed.

Keywords: Broiler finisher, Phytase, Roxazyme G, Growth, carcass characteristics

INTRODUCTION

Broiler, a meat-type of poultry, has the ability to grow fast and reach market weight faster than ruminants (Madubuike and Ekenyem, 2001) and has stirred up interest in many farmers, because it plays a significant role as animal protein source in human diet. Cereal grains supply the bulk of livestock feed, especially for poultry and pigs. However, in developing countries like Nigeria, cereal grains are in high demand for human uses and the production is inadequate to meet the need of the increasing population, consequently, there is little or no excess grain for livestock feed and when available, it is always very expensive. Agro-industrial by-products (AIBs) are important feed components in poultry diets in Nigeria mainly due to the increased competition for the conventional ingredients by humans and the food industries (Iyayi and Davies, 2005). Those high fibre contents are being used either as fillers or as energy diluents or both. For example maize offal, rice offal (Tuleun *et al.*, 2009), brewer's dry grain (Iyayi and Davies, 2005) and palm kernel meal (Ezieshi and Olomu, 2004) have been employed in the formulation of poultry feeds. However, inclusion of high levels of some of the AIBs or the use of high-fibre containing

ones in poultry diets is limited due to their effect of reduced performance in birds (Iyayi and Davies, 2005), reduced nutrient utilization and precipitate metabolic dysfunction with attendant growth depression when ingested by non-ruminants (Onyimonyi, 2005). To enhance the nutritive value of these agro-industrial by-products poultry farmers result to inclusion of feed additives which include antimicrobials, antioxidants, emulsifiers, binders, acidifiers and enzymes. Phytic acid, *myo*-inositol hexaphosphate, is present in grains and oilseeds and is most known for chelating divalent cations (Deshpande and Cheryan, 1984). Since phytic acid and its salts (phytates) usually occur simultaneously in many seeds, researchers often refer to them as "phytates". Phytate in plant seeds is predominantly present as the calcium-magnesium-potassium salt called "phytin" (Swick and Ivey, 1992). The amount of phytate varies from 0.06-2.22% in cereals and 0.08-6% in cereal-milled fractions and protein products (Reddy, 2002). Phytates readily form insoluble complexes with divalent cations in weak acidic to neutral pH conditions and, consequently, reduces their bioavailability in animals. It also reacts directly or indirectly with charged groups of proteins,

amino acids, or form hydrogen bonds with starch, which may result in an adverse influence on protein and starch digestibility (Swick and Ivey, 1992; Maenz, 2001). Dietary phytase; an esterase that hydrolyzes phytic acid is of nutritional importance because the mineral binding strength decreases and the solubility increase, resulting in an increased bioavailability of essential dietary minerals (P and Ca), energy and amino acids (Cowieson *et al.*, 2006). Roxazyme G is an enzyme complex derived from *Trichoderma viride*. It has glucanase and xylanase activity which hydrolysis non starch polysaccharides into smaller molecules which can be utilized and digested by poultry (Broz and Frigg, 1990). This study was therefore carried out to determine the response of broiler finishers to diets supplemented with Roxazyme G and exogenous phytase.

MATERIALS AND METHODS

The experiment was carried out at the Poultry Unit, Teaching and Research Farm, University of Ibadan, Nigeria, between October and December. Roxazyme G and phytase used in the study were purchased from a commercial farm product store in Ibadan, Oyo state, Nigeria.

Experimental birds and diets

A total of 480 of 4-week old unsexed Abor Acre broilers of body weights between 779.58 - 783.33g were randomly allotted to four dietary treatments.

Each treatment was replicated 3 times with 40 birds per replicate. All necessary medications and vaccinations were carried out as recommended by the breeder. The birds were brooded for 3 weeks on a standard broiler starter feed and water was supplied *ad-libitum*. Diet 1 (control) was formulated to meet nutritional requirements of the birds at finisher phase without enzyme supplementation. Diet 2 contained Roxazyme G at 0.2g/kg; Diet 3 contained Phytase at 1g/kg while Diet 4 contained Roxazyme G and Phytase at 0.2g/kg and 1g/kg respectively. The gross composition of starter and finisher diets are shown in Table 1

Data collection

At the beginning of the experiment, the birds were weighed and subsequently weighed weekly. Feed intake was recorded daily; feed conversion ratio was calculated as quantity of feed consumed per unit weight gained over the same period. Mortality during the duration of the study was also recorded. At the end of the experiment, three birds per replicate whose weight Table 1: Gross composition of starter and finisher diets were close to the mean weight of 1.9kg were

Ingredients (%)	Starter	Finisher
Maize	54.13	55.00
Soyabean meal	30.00	21.00
Groundnut cake	5.00	10.00
Wheat offal	0.00	5.88
Fish meal	3.00	0.00
Palm oil	2.00	2.00
Oyster shell	2.00	3.00
Dicalcium phosphate	2.50	2.00
Premix*	0.25	0.25
L- lysine	0.10	0.20
DL-Methionine	0.20	0.20
Table salt	0.25	0.25
Mycofix	0.02	0.02
Biotronics	0.05	0.20
TOTAL	100	100
Calculated value		
Crude protein (%)	22.53	19.70
Energy (ME kcal/kg)	3010.4	3124.0
Crude fibre (%)	3.31	3.72
Calcium (%)	1.44	0.90
Available P. (%)	0.68	0.45
Methionine	0.55	0.49
Lysine	1.30	1.11

Premix* provided the following per kg of diet vitamin A 12500 I.U. Vit D₃ 2500 I.U. vit E 50 mg vit K₃ 2.5mg; vit B₁ 3.0mg; vit B₂ 6.0mg; vit B₆ 6.0mg; niacin 40.0mg; calcium pantothenate 10mg; Biotin 0.80mg; vit B₁₂ 0.25mg; folic acid 1.0mg; choline chloride 300mg; manganese 100mg; iron 50mg; zinc 45mg; cobalt 0.25mg; iodine 1.55mg; selenium 0.1mg.

slaughtered and carcass measurement was taken. Cost/kg of the diet, cost/kg weight gain of bird and cost of total feed consumed were also calculated.

Data analysis

All data collected were subjected to one-way Analysis of variance using SAS (version 9.2) package and means were separated using Duncan multiple range test of the same software.

RESULTS AND DISCUSSION

The results of performance characteristics of broiler finishers fed enzyme supplemented diets are presented in Table 2. There were no significant differences ($p>0.05$) among treatments in average daily weight gain (ADWG), average daily feed intake (ADFI) and feed conversion ratio (FCR). Birds fed on enzyme supplemented diets have been reported to have decreased feed intake as a result of fulfilling their nutrient requirement by taking less amount of feed (Samarasinghe *et al.*, 2000; Oladunjoye and Ojebiyi, 2010). However, the studies of Manafi *et al.* (2011)

Table 2: Performance characteristics of broiler finishers fed diets supplemented with Roxazyme G and Phytase (5 to 8 weeks)

Parameters	T1	T2	T3	T4	SEM(±)
Initial body weight (g)	782.5	783.3	779.6	781.3	2.54
Final body weight (g)	1974.4	1972.1	1888.7	1889.1	24.14
Average daily feed intake (g)	137.9	137.7	136.4	136.1	0.59
Average weight gain (g)	1191.9	1188.8	1109.2	1107.9	22.61
Average daily weight gain (g)	59.6	59.4	55.5	55.4	1.13
Feed conversion ratio	1.4	1.4	1.6	1.4	0.02
Mortality (%)	3.00	1.00	2.00	1.00	0.41

^{abc} Means with same superscript are not significantly different ($p > 0.05$)

T1: Basal diet (without enzyme inclusion)

T2: Basal diet with Roxazyme G inclusion at 0.2g/kg

T3: Basal diet with Phytase inclusion at 1g/kg

T4: Basal diet with Roxazyme G and Phytase at 0.2g/kg and 1g/kg respectively

suggested that broiler finishers fed enzyme supplemented diets have increased feed intake due to increased nutrient digestibility. This however contradicted the findings of this study. This may be as a result of the age of the birds used in this study as broilers in the finishing phase utilise feed nutrient more efficiently (Iyayi and Adegboyega, 2004). The results of the carcass yield and cut up parts as influenced by enzyme supplementation is presented in Table 3. There were observed significant ($p < 0.05$) differences in hot carcass weight and dressing percentage among dietary treatments. Carcass weight of birds on dietary T4 was significantly ($p < 0.05$) different from birds on dietary T3.

However, the dressing yield (%) also showed significant ($p < 0.05$) increase across dietary treatment with birds on T4 showing significant ($p < 0.05$) difference from birds on other dietary treatment. Alam *et al.* (2003) and Rahman *et al.* (2005) had reported that enzyme addition in diet increases fat deposition in carcass and breast muscles and a resultant increase in carcass yield. However, the result of this study is contradicted by the findings of Biswas *et al.* (1999) and Nwoche *et al.* (2006). They reported that carcass yield did not differ among enzyme supplemented diets in broiler finishers. The cut up parts results of neck, shank, thigh, wings and back meat were not significantly ($p > 0.05$) different across all dietary treatments. This showed that dietary enzymes inclusion at broiler finisher phase had no effect on tissue synthesis. These results support the findings of Ahmed *et al.* (2004) who reported that primal cut parts did not differ significantly ($p > 0.05$) but were significant at $p < 0.01$. This was also corroborated by the findings of Scheideler and Ferket (2000), Alam *et al.* (2003). Weights of the head,

drumstick and breast meat though were significantly ($p < 0.05$) different fell within the levels reported by Isikwenu *et al.* (2010). These values were highest in enzyme supplemented diets supported by the findings of Leeson *et al.* (1996), Ahmed *et al.* (2004), Rahman *et al.* (2005), except the drumstick weight which was highest in the control diet in contradiction to the findings of Scheideler and Ferket (2000), Alam *et al.* (2003). Scheideler and Ferket (2000) suggested that improvements in drumstick and thigh weights of enzyme supplemented diet was probably as a result of increase in bone ash and proposed improved tibial bone strength as muscle development was enhanced by improved mobility. However, no differences ($p > 0.05$) was observed in breast meat weights of the same study. The breast muscles and drumsticks are the most economically important portion of the carcass and also provide the greatest portions of edible meat in broilers. Table 4 shows the gastrointestinal characteristics of broiler finishers as influenced by enzyme supplementation. There were no significant ($p > 0.05$) differences in the intestinal length and weight across the dietary treatment. This supported the findings of Wang (2000), Iji *et al.* (2001), Awad *et al.* (2009). They concluded that this was due to reduction of the viscous environment in the intestinal lumen of birds caused by water-soluble NSP. The length of the ceca and weight of ceca and ileum also showed no significant ($p > 0.05$) increase. The ileum weight is an indication of the rate of absorption of nutrient. This finding may be an indication that enzyme supplementation had no positive influence on the absorption of nutrients in broiler finishers used in this study. There were no significant ($p > 0.05$) differences in the weight of the full and empty proventriculus.

Table 3: Cut up parts (% of live weight) of broiler finishers fed diets supplemented with Roxazyme G and phytase.

Parameters	T1	T2	T3	T4	SEM(±)
Head	2.48 ^b	2.87 ^{ab}	2.59 ^{ab}	3.17 ^a	0.11
Neck	4.42	4.61	4.61	5.16	0.18
Shank	4.08	3.81	3.69	4.08	0.11
Drumsticks	11.52 ^a	10.31 ^{ab}	9.74 ^{ab}	8.62 ^b	0.33
Thigh	11.05	10.36	10.96	10.57	0.26
Wings	8.38	8.18	7.92	8.27	0.10
Breast	13.23 ^{ab}	12.21 ^b	12.62 ^b	15.03 ^a	0.40
Back	12.54	12.16	12.30	12.75	0.14
Carcass weight (g)	1426.67 ^{ab}	1467.78 ^{ab}	1395.56 ^b	1533.33 ^a	21.11
Dressing percentage (%)	76.89 ^b	72.68 ^c	71.96 ^c	80.78 ^a	0.83

^{ab} Means with same superscript are not significantly different ($p > 0.05$)

Table 4: Gastrointestinal characteristics of broiler finishers fed diets supplemented with Roxazyme G and phytase

Parameters	T1	T2	T3	T4	SEM(±)
Full proventriculus (g)	8.71	9.64	8.75	9.59	0.24
Empty proventriculus (g)	8.09	8.86	8.29	8.98	0.21
Full crop (g)	13.02 ^a	11.18 ^{ab}	10.24 ^b	9.60 ^b	0.47
GIT weight (g)	102.96	108.19	91.30	96.79	3.35
Intestinal length (cm)	222.13	241.77	218.91	239.33	4.52
Weight of 2 cm ileum (g)	0.67	0.57	0.63	0.51	0.03
Ceca weight (g)	13.51	12.94	11.56	11.29	0.56
Ceca length (cm)	41.65	41.33	39.33	42.64	0.07

^{ab} Means with same superscript are not significantly different ($p > 0.05$)

T1: Basal diet (without enzyme inclusion)

T2: Basal diet with Roxazyme G inclusion at 0.2g/kg

T3: Basal diet with Phytase inclusion at 1g/kg
respectively

T4: Basal diet with Roxazyme G and Phytase at 0.2g/kg and 1g/kg

Table 5: Cost analyses of broiler finishers fed diets supplemented with Roxazyme G and phytase

Parameters	T1	T2	T3	T4
Cost/kg diet (₦)	88.3	90.8	89.3	91.8
Cost of feed/kg weight gain (₦)	204.3	210.4	219.58	225.5
Cost of total feed consumed (₦)	243.8	249.8	243.76	249.7

T1: Basal diet (without enzyme inclusion).

T2: Basal diet with Roxazyme G inclusion at 0.2g/kg.

T3: Basal diet with Phytase inclusion at 1g/kg.
respectively

T4: Basal diet with Roxazyme G and Phytase at 0.2g/kg and 1g/kg

Wang (2000) concluded that an inclusion of enzyme preparation high in xylanase activity in a diet caused decreased weight of proventriculus. However, there was significant difference in weight of full crop. The value was highest in birds on control diet which probably indicates that enzyme inclusion in diet increases rate at which feed is digested in the GIT. The results of cost analysis are presented in Table 5. The result of this study shows that enzyme inclusion in an iso-caloric and iso-nitrogenous broiler finisher diet marginally increased the cost of feed/kg and consequently raises cost of production. Also the cost of feed/kg weight gain by bird also increased across the dietary treatments. This contradicted the result of Alam *et al.* (2003) and Rahman *et al.* (2005). They reported that enzyme inclusion in an iso-caloric and iso-nitrogenous broiler finisher diet reduced the feed cost/ kg weight gain

thereby, increasing profit. This contradiction is probably because birds on control diet had the highest average weight gain in the course of the study. The cost of total feed consumed by bird also showed similar levels across dietary treatments. This may be explained by the average weight gain of birds on enzyme supplemented diets which is lower than the control diet. This supported the report of Alam *et al.* (2003) that concluded profitability may differ among enzymes signifying the importance of proper selection of enzyme to get best result.

CONCLUSION

The results obtained from this study indicated no significant increase in the performance, carcass yield, internal organ characteristics of broiler finishers fed diets supplemented with Roxazyme G and exogenous

Phytase. The cost analysis also indicated that it is not profitable to include enzymes in an iso-caloric and iso-nitrogenous broiler finishers' diet as this placed an additional cost on production. It is therefore concluded that, inclusion of enzymes in diet of broiler finishers on the same caloric and protein content of nutrient concentration did not improve performance, carcass yield and cost benefit.

CONFLICT OF INTEREST

Authors declare that there is no conflict of interest concerning this manuscript.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the Teaching and Research Farm, University of Ibadan for the provision of the birds and feed for this research.

REFERENCES

- Acamovic T, 2001. Commercial application of enzyme technology for poultry production. *World's Poultry Science Journal* 57: 225-242.
- Ahmed, F., Rahman, M. S., Ahmed, S. U. and Miah, M.Y. 2004. Performance of broiler supplemented soybean meal based diet. *International Journal of Poultry Science*, 3(4): 266-271
- Alam, M. J., Howlider, M. A. R., Pramanik, M. A. H. and Haque, M. A. 2003. Effect of Exogenous Enzyme in Diet on Broiler Performance. *International Journal of Poultry Science*, 2: 168-173.
- Awad, W. A., Ghareeb, K., Abdel-Raheem, S. and Bohm, J. 2009. Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights, and intestinal histomorphology of broiler chickens. *Poultry Science*, 88, 49-55.
- Biswas T, Mandal L and Sarker S. K. 1999. Studies of enzymes supplementation and herbal preparation at different levels of energy on the performance of broilers. *Journal of Interacademic*, 3(1): 53-58.
- Broz, J. and Frigg, R.I. 1990. Influence of *Trichoderma viride* enzyme complex on nutrient value of barley and oats for broilers. *Archive*, 54:34-37.
- Cowieson, A. J., Acamovic, T. and Bedford, M. R. 2006. Supplementation of corn-soy based diets with *Escherichia coli*-derived phytase: effects on broiler chick performance and the digestibility of amino acids and metabolizability of minerals and energy. *Poultry Science*. 85:1389-1397.
- Deshpande, S.S. and Cheryan, M. 1984. Effects of phytic acid, divalent cations, and their interaction on α -amylase activity. *Journal of Food Science*. 49:516-519.
- Ezieshi, E.V. and Olomu, J. M. 2004. Comparative performance of broiler chickens fed varying levels of Palm Kernel cake and maize offal. *Pakistan Journal of Nutrition*. 3: 254-257.
- Iji, P.A., Saki, A.A. and Tivey, D.R. 2001. Intestinal structure and function of broiler chickens on diets supplemented with a mannan oligosaccharide. *Journal of Science and Food Agriculture* 81, 1186-1192.
- Isikwenu, J. O, Akpodiete, O. J., Omeje, S. I. and Okagbare, G. O. 2010. The effect of replacing groundnut cake and urea treated and fermented brewer's dried grains on nutrient digestibility, retention and carcass characteristic of broiler finishers. *Nigerian Journal of Animal Production*. 37: 1-12.
- Iyayi, E. A. and Adegboyega, B. A. 2004. Supplementation of wheat bran and brewer's dried grain diets with Roxazyme G enzyme for broiler feeding. In Proceedings of 2004 BSAS Annual Conference. University of York, York, UK, April 5-7. P 137.
- Iyayi, E.A. and Davies, B. I. 2005. Effect of Enzyme supplementation of Palm Kernel meal and Brewer's Dried Grain on the performance of broilers. *International Journal of Poultry Science* 4 (2): 76-80.
- Lesson S, Caston, L J and Yublut, D 1996. Adding Roxazyme to wheat diets of chickens and turkey broilers; *Journal of Poultry Research*. 5(2): 167-172.
- Madubuike, F.N. and Ekenyem, B.U. 2001. Non-ruminant livestock production in the tropics. Gust graphics centre, Owerri, Nigeria. Pp 120-145.
- Maenz, D. D. 2001. Enzymatic characteristics of phytases as they relate to their use in animal feeds. In: Enzymes in Farm Animal Nutrition (M. R. Bedford and G. G. Partridge, Eds.), CABI Publishing, New York, NY. pp. 61-84.
- Manafi, M., Hossein, B. and Mohammad, Y. 2011. Effect of Polyzyme in Broilers Fed with Corn (*Zea mays* L.) Bran-Based Diets. *Advances in Environmental Biology*, 5(7): 1651-1655.
- Nwoche, G.N., Ndubuisi, E.C. and Iheukwumere, F.C. 2006. Performance of finisher broilers and cost implication of feeding palm oil as energy supplement. *Nigerian Agriculture Journal*. 37: 44-49.
- Oladunjoye, I.O. and Ojebiyi, O.O. 2010. Performance Characteristics of Broiler Chicken (*Gallus gallus*) Fed Rice (*Oryza sativa*) Bran with or Without Roxazyme G2G. *International Journal of Animal and Veterinary Advances* 2 (4): 135-140.

- Onyimonyi, A.E. 2005. Comparative evaluation of the performance of laying birds fed Safizyme GP 800®. *Proceedings of 30th Annual Conference of Nigeria Society for Animal Production*. Held at Nsukka on March 2005. Pp 228-230.
- Rahman, M. M., Mollah, M. B. R., Islam, F. B. and Howlider, M. A. R. 2005. Effect of enzyme supplementation to parboiled rice polish based diet on broiler performance. *Livestock Research for Rural Development*. 17(4): 21-24.
- Reddy, N. R. 2002. Occurrence, distribution, content, and dietary intake of phytate. *In: Food Phytates* (N. R. Reddy and S. K. Sathe, Eds.), CRC Press LLC, Boca Raton, FL. pp. 25-51.
- Samarasinghe, K., Messikommer, R. and Wenk, C. 2000. Activity of supplemental enzymes and their effect on nutrient utilization and growth performance of growing chickens as affected by pelleting temperature. *Archives of Animal Nutrition* 53, 45–58.
- Scheideler, S. E. and Ferket, P. R. 2000. Phytase in broiler rations- Effects on carcass yields and Incidence of tibial dyschondroplasia. *Poultry Science Association, Inc.* 468-475.
- Swick, R.A. and Ivey, R.J. 1992. The value of improving phosphorus retention. *Feed Management*. 43:8-17.
- Tuleun, C.D., Yaakugh, I.D.I. and Okwori, A.I. 2009. Performance of pullets fed on graded levels of rice offal supplemented with Roxazyme G® enzyme. *Journal of Applied Biosciences* 20: 1153 – 1158
- Wang, G. J. 2000. Effects of enzyme supplementation and irradiation of rice bran on the performance of Leghorn and broiler chicks. *In: Enzymes in Poultry and Swine Nutrition* (R. R. Marquardt and Z. Han Eds.), International Development Research Centre, Ottawa, Canada.